arXiv:1503.02116v1 [math.RT] 7 Mar 2015

FINITENESS OF THE NUMBER OF MINIMAL ATOMS IN GROTHENDIECK CATEGORIES

RYO KANDA

Abstract. For a Grothendieck category having a noetherian generator, we prove that there exist only finitely many minimal atoms. This is a far-reaching generalization of the well-known fact that every commutative noetherian ring has only finitely many minimal prime ideals. It is also shown that each minimal atom is represented by a compressible object.

1. Introduction

Gabriel's spectrum [Gab62] is a fundamental tool to study Grothendieck categories and has been investigated by many authors ([Her97, Kra97, Ros95, Pap02] for example). The *atom spectrum* of a Grothendieck category is a variant of Gabriel's spectrum, which was introduced by the author [Kan12] based on the work of [Sto72]. Each element of the atom spectrum is called an *atom*. While Gabriel's spectrum is defined as the set of isomorphism classes of indecomposable injective objects, the atom spectrum is defined as the set of equivalence classes of monoform objects in order that it works even for other kinds of abelian categories without enough injective objects such as noetherian abelian categories (see [Kan12]). Moreover, the atom spectrum always has the structure of a partially ordered set, and in the case of the category of modules over an arbitrary commutative ring, it is isomorphic to the prime spectrum of the commutative ring as a partially ordered set.

In the case of locally noetherian Grothendieck categories, such as the category $\operatorname{Mod} \Lambda$ of right modules over a right noetherian ring Λ, the atom spectrum coincides with Gabriel's spectrum via generalized Matlis' correspondence ([Kan12, Theorem 5.9]) and behaves as the set of structural elements of the Grothendieck category. For example, the atom spectrum is large enough to classify localizing subcategories (Theorem 3.3) and E-stable subcategories ([Kan15, Theorem 6.4]).

In this paper, we investigate a Grothendieck category having a noetherian generator and its atom spectrum. The category of right modules over an arbitrary right noetherian ring is our motivating example. The advantage of this approach is that a Grothendieck category having a noetherian generator can be localized at *every* atom and we obtain such a Grothendieck category again as the output. This fact enables us to show new results which have not been revealed in classical ring theory.

For a Grothendieck category with a noetherian generator, our previous result states that there exist enough minimal atoms in the sense that each atom is greater than some minimal atom (Proposition 4.2). As Theorem 4.4, we prove the main result of this paper:

Theorem 1.1. *Let $\mathcal{A}$ be a Grothendieck category having a noetherian generator. Then there exist only finitely many minimal atoms in $\mathcal{A}$.*

This is a far-reaching generalization of the well-known fact that every commutative noetherian ring has only finitely many minimal prime ideals. Since the minimal atoms are not necessarily *associated atoms* of the fixed noetherian generator (Remark 4.8), we shall develop another method using the notion of compressible objects, which has been studied in [Gol64, Gol72, Mew72, GR73,

---

2010 *Mathematics Subject Classification.* 18E15 (Primary), 16P40, 16D90, 14A22, 13C60 (Secondary).

Key words and phrases. Minimal atom; Grothendieck category; noetherian generator; compressible object.

The author is a Research Fellow of Japan Society for the Promotion of Science. This work is supported by Grant-in-Aid for JSPS Fellows 25·249.

Goo80] for example. A key step in the proof is given by Theorem 4.1 asserting that every minimal atom is represented by a compressible object.

The main theorem yields a new result even for right noetherian rings.

Corollary 1.2. *Let Λ be a right noetherian ring. Then there exist only finitely many minimal atoms in* $\operatorname{Mod}\Lambda$.

2. Acknowledgement

The author would like to express his deep gratitude to Osamu Iyama for his elaborated guidance.

3. Preliminaries

This section is devoted to preliminary results which we use in the proof of the main result. We refer the reader to [Pop73] for the general theory of abelian categories and Grothendieck categories.

Throughout this section, let $\mathcal{A}$ be a Grothendieck category. We recall the definition of the atom spectrum of $\mathcal{A}$ and some basic results. See [Kan14, section 3] for more details.

Definition 3.1.

(1) A nonzero object H in $\mathcal{A}$ is called *monoform* if for each nonzero subobject L of H, no nonzero subobject of H is isomorphic to a subobject of H/L.

(2) For monoform objects H_1 and H_2 in $\mathcal{A}$, we say that H_1 is *atom-equivalent* to H_2 if there exists a nonzero subobject of H_1 which is isomorphic to a subobject of H_2. This is an equivalence relation on the class of monoform objects in $\mathcal{A}$ ([Kan12, Proposition 2.8]).

(3) The *atom spectrum* $\operatorname{ASpec}\mathcal{A}$ of $\mathcal{A}$ is the quotient set of the class of monoform objects in $\mathcal{A}$ by the atom equivalence. Each element of $\operatorname{ASpec}\mathcal{A}$ is called an *atom* in $\mathcal{A}$. For each monoform object H in $\mathcal{A}$, the equivalence class of H is denoted by $\overline{H}$.

For a ring Λ, denote by $\operatorname{Mod}\Lambda$ the category of right Λ-modules. This is a Grothendieck category. For a commutative ring R, we have a bijection between the prime spectrum $\operatorname{Spec} R$ and the atom spectrum $\operatorname{ASpec}(\operatorname{Mod} R)$ given by $\mathfrak{p} \mapsto \overline{R/\mathfrak{p}}$ (see [Sto72, p. 631] or [Kan12, Proposition 7.2 (1)]). Moreover, the notion of support in commutative ring theory is generalized to Grothendieck categories as follows.

Definition 3.2. Let M be an object in $\mathcal{A}$. The *atom support* $\operatorname{ASupp} M$ of M is the subset of $\operatorname{ASpec}\mathcal{A}$ defined by

$$\operatorname{ASupp} M = \{\, \alpha \in \operatorname{ASpec}\mathcal{A} \mid \alpha = \overline{H} \text{ for some monoform subquotient } H \text{ of } M \,\}.$$

The atom spectrum $\operatorname{ASpec}\mathcal{A}$ is regarded as a topological space with respect to the *localizing topology*, that is, a subset Φ of $\operatorname{ASpec}\mathcal{A}$ is open if and only if $\Phi = \operatorname{ASupp} M$ for some object M in $\mathcal{A}$. For a commutative ring R, a subset of $\operatorname{ASpec}(\operatorname{Mod} R)$ is open if and only if the corresponding subset Φ of $\operatorname{Spec} R$ is closed under specialization, that is, the conditions $\mathfrak{p} \in \Phi$ and $\mathfrak{p} \subset \mathfrak{q}$ imply $\mathfrak{q} \in \Phi$. The localizing topology is used to classify the localizing subcategories of $\mathcal{A}$:

Theorem 3.3 ([Her97, Theorem 3.8], [Kra97, Corollary 4.3], and [Kan12, Theorem 5.5])**.** *Let $\mathcal{A}$ be a locally noetherian Grothendieck category. Then we have a bijection*

$$\{\, \textit{localizing subcategories of } \mathcal{A} \,\} \to \{\, \textit{open subsets of } \operatorname{ASpec}\mathcal{A} \,\}$$

given by $\mathcal{X} \mapsto \operatorname{ASupp}\mathcal{X} := \bigcup_{M \in \mathcal{X}} \operatorname{ASupp} M$. The inverse map is given by

$$\Phi \mapsto \operatorname{ASupp}^{-1}\Phi := \{\, M \in \mathcal{A} \mid \operatorname{ASupp} M \subset \Phi \,\}.$$

Proof. See [Kan14, Theorem 6.8] and [Kan14, Proposition 6.5] for a detailed proof. □

The inclusion relation between prime ideals is generalized as the specialization order on $\operatorname{ASpec}\mathcal{A}$, which is denoted by $\leq$ (see [Kan14, Definition 3.16]).

For a localizing subcategory $\mathcal{X}$, the quotient category $\mathcal{A}/\mathcal{X}$ is a Grothendieck category together with a canonical (covariant) functor $\mathcal{A} \to \mathcal{A}/\mathcal{X}$, which is dense and exact and has a fully faithful right adjoint (see [Kan13, section 5]). For each $\alpha \in \operatorname{ASpec}\mathcal{A}$, the localization $\mathcal{A}_\alpha$ is defined as the quotient category of $\mathcal{A}$ by the localizing subcategory corresponding to α ([Kan13, Definition 6.1]), and the canonical functor $\mathcal{A} \to \mathcal{A}_\alpha$ is denoted by $(-)_\alpha$. This is a generalization of the localization of a commutative ring at a prime ideal ([Kan13, Proposition 6.9]).

The following result is used to obtain a subobject in $\mathcal{A}$ from a subobject in the localized category.

Proposition 3.4. *Let M be an object in $\mathcal{A}$ and $\alpha \in \operatorname{ASpec}\mathcal{A}$. Then for each subobject L' of M_α, there exists a subobject L of M such that*

1. *$L_\alpha = L'$,*
2. *L is largest with respect to the property $L_\alpha \subset L'$, and*
3. *$N_\alpha \neq 0$ for every nonzero subobject N of M/L.*

Proof. This follows from [Kan14, Proposition 4.9] and [Kan13, Proposition 6.2]. □

We introduce a characterization of monoform objects. Recall that every nonzero subobject of a monoform object is again monoform ([Kan12, Proposition 2.2]).

Proposition 3.5. *Let H be an object in $\mathcal{A}$ and $\alpha \in \operatorname{ASpec}\mathcal{A}$. Then H is a monoform object with $\overline{H} = \alpha$ if and only if*

1. *H_α is a simple object in $\mathcal{A}_\alpha$, and*
2. *$L_\alpha \neq 0$ for every nonzero subobject L of H.*

Proof. Assume that H is a monoform object with $\overline{H} = \alpha$. For every nonzero subobject L of H, we have $\alpha = \overline{H} = \overline{L} \in \operatorname{ASupp} L$, and hence $L_\alpha \neq 0$ by [Kan13, Proposition 6.2]. Let N' be a nonzero subobject of H_α. Then by Proposition 3.4, there exists a subobject N of H satisfying $N_\alpha = N'$. By [Kan13, Proposition 2.14], we have $\alpha \notin \operatorname{ASupp}(H/N)$, and hence $H_\alpha/N_\alpha \cong (H/N)_\alpha = 0$. This shows that $N' = N_\alpha = H_\alpha$, and H_α is simple.

Assume that H satisfies the latter two conditions. Let L be a nonzero subobject of H. By the assumption, we have $(H/L)_\alpha \cong H_\alpha/L_\alpha = 0$, and hence $N_\alpha = 0$ for every subobject N of H/L. Therefore no nonzero subobject of H is isomorphic to a subobject of H/L. This shows that H is monoform. Since we have $\alpha \in \operatorname{ASupp} H$, there exist subobjects $L_1 \subset L_2 \subset H$ such that L_2/L_1 is monoform and $\overline{L_2/L_1} = \alpha$. If $L_1 \neq 0$, then $(L_2/L_1)_\alpha \subset (H/L_1)_\alpha = 0$ by the above argument. This contradicts $\alpha = \overline{L_2/L_1} \in \operatorname{ASupp}(L_2/L_1)$. Hence $L_1 = 0$, and $\overline{H} = \overline{L_2} = \alpha$. □

Recall that a *generator* in a Grothendieck category $\mathcal{A}$ is an object M such that every object in $\mathcal{A}$ is a quotient object of a (possibly infinite) direct sum of copies of M. In the next section, we investigate a Grothendieck category having a noetherian generator. The following well-known result shows that this property is preserved by taking the quotient category by a localizing subcategory.

Proposition 3.6. *Let $\mathcal{X}$ be a localizing subcategory of $\mathcal{A}$, and let $F\colon \mathcal{A} \to \mathcal{A}/\mathcal{X}$ be the canonical functor.*

1. *If M is a generator in $\mathcal{A}$, then $F(M)$ is a generator in $\mathcal{A}/\mathcal{X}$.*
2. *For every noetherian object M in $\mathcal{A}$, the object $F(M)$ in $\mathcal{A}/\mathcal{X}$ is noetherian.*

Proof. (1) This is shown by using the fact that F is dense and exact and preserves arbitrary direct sums.

(2) This follows from [Kan14, Proposition 4.9]. □

The following lemmas are used in the proof of the main result in the next section.

Lemma 3.7. *If M is a generator in $\mathcal{A}$, then* $\operatorname{ASpec}\mathcal{A} = \operatorname{ASupp} M$.

Proof. Let H be a monoform object in $\mathcal{A}$. Then there exists a nonzero morphism $f\colon M \to H$. Since $\operatorname{Im} f$ is a nonzero subobject of H, we have $\overline{H} = \overline{\operatorname{Im} f} \in \operatorname{ASupp}(\operatorname{Im} f) \subset \operatorname{ASupp} M$. Hence the assertion holds. ☐

Lemma 3.8. *Let $\alpha \in \operatorname{ASpec} \mathcal{A}$.*

1. *Let H be a monoform object in $\mathcal{A}$ with $\overline{H} = \alpha$, and let M be an object in $\mathcal{A}$ satisfying $M_\alpha = 0$. Then $\operatorname{Hom}_{\mathcal{A}}(M, H) = 0$.*
2. *Let H_1 and H_2 be monoform objects with $\overline{H_1} = \overline{H_2} = \alpha$. Then every nonzero morphism $H_1 \to H_2$ is a monomorphism.*

Proof. (1) Assume that a morphism $f\colon M \to H$ is nonzero. Then we have $\alpha = \overline{H} = \overline{\operatorname{Im} f} \in \operatorname{ASupp}(\operatorname{Im} f)$. On the other hand, since $\operatorname{Im} f$ is a quotient object of M, we have $(\operatorname{Im} f)_\alpha = 0$ by the assumption. By [Kan13, Proposition 6.2], we have $\alpha \notin \operatorname{ASupp}(\operatorname{Im} f)$. This is a contradiction.

(2) Assume that a morphism $f\colon H_1 \to H_2$ is not a monomorphism. Then Proposition 3.5 shows that $(\operatorname{Ker} f)_\alpha$ is a nonzero subobject of the simple object $(H_1)_\alpha$, and we have $(H_1/\operatorname{Ker} f)_\alpha \cong (H_1)_\alpha/(\operatorname{Ker} f)_\alpha = 0$. By (1), the induced monomorphism $H_1/\operatorname{Ker} f \to H_2$ is zero, and we obtain $f = 0$. ☐

The notion of compressible object, which has been studied in [Gol64, Gol72, Mew72, GR73, Goo80] for example, plays an important role in the next section. We recall the definition and some basic properties of compressible objects.

Definition 3.9. A nonzero object H in $\mathcal{A}$ is called *compressible* if each nonzero subobject L of H has some subobject which is isomorphic to H.

Proposition 3.10.

1. *Every nonzero subobject of a compressible object in $\mathcal{A}$ is compressible.*
2. *If $\mathcal{A}$ is a locally noetherian Grothendieck category, then every compressible object in $\mathcal{A}$ is noetherian and monoform.*

Proof. (1) Straightforward.

(2) Let H be a compressible object in $\mathcal{A}$. Then H has a nonzero noetherian subobject, and by [Kan12, Theorem 2.9], the subobject has a monoform subobject L. There exists a subobject of L which is isomorphic to H, and hence H is monoform. ☐

In the case of commutative rings, compressible modules are characterized as follows.

Proposition 3.11. *Let R be a commutative ring.*

1. ([Gol72, 3.38]) *For an ideal $\mathfrak{p}$ of R, the object $R/\mathfrak{p}$ in $\operatorname{Mod} R$ is compressible if and only if $\mathfrak{p}$ is a prime ideal.*
2. *An R-module H is a compressible object in $\operatorname{Mod} R$ if and only if H is isomorphic to a nonzero R-submodule of $R/\mathfrak{p}$ for some $\mathfrak{p} \in \operatorname{Spec} R$.*

Proof. (1) For the convenience of the reader, we include a proof.

Assume that $R/\mathfrak{p}$ is compressible. Then for each $a \in R \setminus \mathfrak{p}$, we have $\operatorname{Ann}_R(\overline{a} \in R/\mathfrak{p}) = \mathfrak{p}$ since $\overline{a}R$ has an R-submodule isomorphic to $R/\mathfrak{p}$. This shows that $\mathfrak{p}$ is a prime ideal.

Conversely, let $\mathfrak{p}$ is a prime ideal, and let L be a nonzero R-submodule of $R/\mathfrak{p}$. Take a nonzero element $\overline{a} \in L$. Since we have $\operatorname{Ann}_R(\overline{a}) = \mathfrak{p}$, it holds that $\overline{a}R \cong R/\mathfrak{p}$.

(2) Assume that H is compressible. Take a nonzero element $x \in H$. Then H is isomorphic to an R-submodule of xR, which is compressible by Proposition 3.10 (1). We have $xR \cong R/\operatorname{Ann}_R(x)$, and by (1), the ideal $\operatorname{Ann}_R(x)$ is prime.

The converse follows from (1) and Proposition 3.10 (1). ☐

Remark 3.12. For a commutative ring R, each atom in $\operatorname{Mod} R$ is of the form $\overline{R/\mathfrak{p}}$ for some $\mathfrak{p} \in \operatorname{Spec} R$, and $\overline{R/\mathfrak{p}}$ is a compressible object in $\operatorname{Mod} R$.

On the other hand, for a right noetherian ring, an atom is not necessarily represented by a compressible object. Indeed, Goodearl [Goo80] constructed a left and right noetherian ring Λ such that there exists a monoform Λ-module which does not have a compressible Λ-submodule.

4. Main result

Throughout this section, let $\mathcal{A}$ be a Grothendieck category having a noetherian generator. Denote by $\operatorname{AMin}\mathcal{A}$ the subset of $\operatorname{ASpec}\mathcal{A}$ consisting of all minimal atoms. The following theorem shows that the behavior of minimal atoms is similar to atoms in $\operatorname{Mod} R$ for a commutative ring R.

Theorem 4.1. *For every $\alpha \in \operatorname{AMin}\mathcal{A}$, there exists a compressible object H in $\mathcal{A}$ satisfying $\alpha = \overline{H}$.*

Proof. Let M be a noetherian generator in $\mathcal{A}$. By [Kan13, Proposition 6.6 (1)], the topological space $\operatorname{ASpec}\mathcal{A}_\alpha$ is homeomorphic to the singleton $\{\alpha\}$ since α is minimal. Hence by Proposition 3.6 (2) and [Kan13, Proposition 3.7 (3)], the object M_α in $\mathcal{A}_\alpha$ has a composition series

$$0 = L'_0 \subset L'_1 \subset \cdots \subset L'_n = M_\alpha.$$

By using Proposition 3.4, we obtain a series

$$L_0 \subset L_1 \subset \cdots \subset L_n = M$$

such that $(L_i)_\alpha = L'_i$ for each $i = 0, \dots, n$, and $N_\alpha \neq 0$ for every nonzero subobject N of M/L_i. For each $i = 1, \dots, n$, the object $(L_i/L_{i-1})_\alpha \cong L'_i/L'_{i-1}$ is simple, and $N_\alpha \neq 0$ for every nonzero subobject N of L_i/L_{i-1}. Hence by Proposition 3.5, the object $H_i := L_i/L_{i-1}$ is monoform and $\overline{H_i} = \alpha$. Since the monoform objects $H_1, \dots, H_n$ represent the same atom α, there exists a monoform object H in $\mathcal{A}$ such that for each $i = 1, \dots, n$, the object H is isomorphic to some subobject of H_i.

Let H' be a nonzero subobject of H. Since M is a generator in $\mathcal{A}$, there exists a nonzero morphism $M \to H'$. By using the series of M, we obtain a nonzero morphism $L_0 \to H'$ or a nonzero morphism $L_i/L_{i-1} \to H'$ for some $i = 1, \dots, n$. Lemma 3.8 (1) implies that the former case does not occur, and Lemma 3.8 (2) implies that the nonzero morphism $H_i = L_i/L_{i-1} \to H'$ is a monomorphism. Thus H' has a subobject which is isomorphic to H by the definition of H. ☐

The following result shows the existence of minimal atoms, which is used in the proof of Proposition 4.3.

Proposition 4.2. *For each $\alpha \in \operatorname{ASpec}\mathcal{A}$, there exists $\beta \in \operatorname{AMin}\mathcal{A}$ satisfying $\beta \leq \alpha$.*

Proof. The assertion follows from [Kan13, Proposition 4.7] and Lemma 3.7. ☐

In order to prove the finiteness of $\operatorname{AMin}\mathcal{A}$, we show some topological properties of it. Although the statement is quite simple, the proof needs our previous results.

Proposition 4.3. *$\operatorname{AMin}\mathcal{A}$ is a closed subset of $\operatorname{ASpec}\mathcal{A}$, and the induced topology on $\operatorname{AMin}\mathcal{A}$ is discrete.*

Proof. Let $\alpha \in \operatorname{AMin}\mathcal{A}$. We show that the subsets

$$V(\alpha) = \{\, \beta \in \operatorname{ASpec}\mathcal{A} \mid \alpha \leq \beta \,\} \quad \text{and} \quad W(\alpha) = \{\, \beta \in \operatorname{ASpec}\mathcal{A} \mid \alpha < \beta \,\}$$

of $\operatorname{ASpec}\mathcal{A}$ are open.

By Theorem 4.1, there exists a compressible object H in $\mathcal{A}$ satisfying $\alpha = \overline{H}$. It follows from [Kan13, Proposition 4.2] that $V(\alpha) = \operatorname{ASupp} H$, which is an open subset of $\operatorname{ASpec}\mathcal{A}$. Since the minimal atom α is a closed point in $\operatorname{ASpec}\mathcal{A}$, the subset $W(\alpha) = V(\alpha) \setminus \{\alpha\}$ of $\operatorname{ASpec}\mathcal{A}$ is open.

By Proposition 4.2, we obtain the equation

$$\operatorname{AMin}\mathcal{A} = \operatorname{ASpec}\mathcal{A} \setminus \bigcup_{\alpha \in \operatorname{AMin}\mathcal{A}} W(\alpha),$$

which implies that $\operatorname{AMin}\mathcal{A}$ is a closed subset of $\operatorname{ASpec}\mathcal{A}$. The equation

$$V(\alpha) \cap \operatorname{AMin}\mathcal{A} = \{\alpha\}$$

shows that α is an open point in $\operatorname{AMin}\mathcal{A}$. ☐

We are ready to prove the main result of this paper.

Theorem 4.4. *Let $\mathcal{A}$ be a Grothendieck category having a noetherian generator. Then* $\operatorname{AMin}\mathcal{A}$ *is a finite set.*

Proof. By Proposition 4.3, the subset $\varPhi := \operatorname{ASpec}\mathcal{A} \setminus \operatorname{AMin}\mathcal{A}$ of $\operatorname{ASpec}\mathcal{A}$ is open. Let $\mathcal{X} := \operatorname{ASupp}^{-1}\varPhi$. Then by [Kan13, Theorem 5.17] and Theorem 3.3, the topological space $\operatorname{ASpec}(\mathcal{A}/\mathcal{X})$ is homeomorphic to $\operatorname{AMin}\mathcal{A}$. Since it is discrete, by Proposition 3.6 and [Kan13, Proposition 3.7 (3)], the image M' of a noetherian generator in $\mathcal{A}$ by the canonical functor $\mathcal{A} \to \mathcal{A}/\mathcal{X}$ is a generator of finite length in $\mathcal{A}/\mathcal{X}$. Let $S'_1, \dots, S'_n$ be the composition factors of M'. It follows from Lemma 3.7 and [Kan12, Proposition 3.3] that

$$\operatorname{ASpec}\mathcal{A} = \operatorname{ASupp}M' = \operatorname{ASupp}S'_1 \cup \dots \cup \operatorname{ASupp}S'_n = \{\overline{S'_1}, \dots, \overline{S'_n}\}. \qquad \square$$

Theorem 4.4 gives a new result even for right noetherian rings.

Corollary 4.5. *Let $\varLambda$ be a right noetherian ring. Then the set* $\operatorname{AMin}(\operatorname{Mod}\varLambda)$ *is nonempty and finite.*

Proof. This follows from Proposition 4.2 and Theorem 4.4 since $\varLambda$ is a noetherian generator in $\operatorname{Mod}\varLambda$. ☐

For a right noetherian domain $\varLambda$, we can describe the minimal atoms in $\operatorname{Mod}\varLambda$ as follows.

Example 4.6. Let $\varLambda$ be a right noetherian ring which is a domain, that is, for each $a, b \in \varLambda$, the condition $ab = 0$ implies $a = 0$ or $b = 0$. Then the object $\varLambda$ in $\operatorname{Mod}\varLambda$ is compressible since the right $\varLambda$-module $a\varLambda$ is isomorphic to $\varLambda$ for each $a \in \varLambda$. Hence by Proposition 3.10 (2), the object $\varLambda$ in $\operatorname{Mod}\varLambda$ is monoform. Since $\varLambda$ is a generator in $\operatorname{Mod}\varLambda$, it follows from Lemma 3.7 that $\operatorname{ASpec}(\operatorname{Mod}\varLambda) = \operatorname{ASupp}\varLambda$. We deduce from [Kan13, Proposition 4.2] that $\operatorname{AMin}(\operatorname{Mod}\varLambda) = \{\overline{\varLambda}\}$.

Wu [Wu88] gave an example of a Grothendieck category with a noetherian generator which does not satisfy the condition Ab4*, that is, direct product is not exact. In particular, the Grothendieck category is not equivalent to the category of right modules over a ring. We give a description of the minimal atoms of the Grothendieck category.

Example 4.7. For simplicity, let R be a commutative noetherian Gorenstein ring with Krull dimension at least 2. Let $\varPhi := \{\, \mathfrak{p} \in \operatorname{Spec}R \mid \operatorname{ht}\mathfrak{p} \geq 2 \,\}$ and

$$\mathcal{X} := \operatorname{Supp}^{-1}\varPhi = \{\, M \in \operatorname{Mod}R \mid \operatorname{Supp}M \subset \varPhi \,\}.$$

Wu [Wu88] showed that the Grothendieck category $\mathcal{A} := (\operatorname{Mod}R)/\mathcal{X}$ does not satisfy Ab4*, whereas the image of R by the canonical functor $\operatorname{Mod}R \to \mathcal{A}$ is a noetherian generator in $\mathcal{A}$.

In this case, the atom spectrum $\operatorname{ASpec}\mathcal{A}$ is isomorphic to $(\operatorname{Spec}R) \setminus \varPhi = \{\, \mathfrak{p} \in \operatorname{Spec}R \mid \operatorname{ht}\mathfrak{p} \leq 1 \,\}$ as a partially ordered set. Therefore $\operatorname{AMin}\mathcal{A}$ is in bijection with the set of minimal prime ideals of R.

Remark 4.8. For a commutative noetherian ring R, the finiteness of the number of minimal prime ideals can be shown by using the equation

$$\operatorname{Min}(\operatorname{Spec}R) = \operatorname{Min}(\operatorname{Supp}R) = \operatorname{Min}(\operatorname{Ass}R)$$

since $\operatorname{Ass}R$ is a finite set. In the case of a Grothendieck category $\mathcal{A}$ with a noetherian generator M, it holds that $\operatorname{Min}(\operatorname{ASpec}\mathcal{A}) = \operatorname{Min}(\operatorname{ASupp}M)$ by Lemma 3.7, and the set $\operatorname{AAss}M$ of associated atoms of M is finite ([Kan12, Remark 3.6]). However, the equality $\operatorname{Min}(\operatorname{ASupp}M) = \operatorname{Min}(\operatorname{AAss}M)$ is far from being true even in the case where $\mathcal{A} = \operatorname{Mod}\varLambda$ for a ring $\varLambda$. For example, let $\varLambda$ be the ring

$$\begin{bmatrix} K & 0 \\ K & K \end{bmatrix}$$

of 2 × 2 lower triangular matrices over a field K. Denote the indecomposable right $\varLambda$-modules by

$$S_1 := \begin{bmatrix} K & 0 \end{bmatrix},\ S_2 := \begin{bmatrix} K & K \end{bmatrix}/\begin{bmatrix} K & 0 \end{bmatrix},\ P_2 := \begin{bmatrix} K & K \end{bmatrix}.$$

Then $\varLambda = S_1 \oplus P_2$ as a right $\varLambda$-module, and we have

$$\operatorname{AMin}(\operatorname{Mod}\varLambda) = \operatorname{Min}(\operatorname{ASupp}\varLambda) = \operatorname{ASupp}\varLambda = \{\overline{S_1}, \overline{S_2}\} \neq \{\overline{S_1}\} = \operatorname{AAss}\varLambda = \operatorname{Min}(\operatorname{AAss}\varLambda).$$

References

[Gab62] P. Gabriel, Des catégories abéliennes, *Bull. Soc. Math. France* **90** (1962), 323–448.

[Gol64] A. W. Goldie, Torsion-free modules and rings, *J. Algebra* **1** (1964), 268–287.

[Gol72] A. W. Goldie, The structure of Noetherian rings, *Lectures on rings and modules (Tulane Univ. Ring and Operator Theory Year, 1970–1971, Vol. I)*, pp. 213–321, Lecture Notes in Math., Vol. 246, *Springer-Verlag, Berlin-Heidelberg-New York*, 1972.

[Goo80] K. R. Goodearl, Incompressible critical modules, *Comm. Algebra* **8** (1980), no. 19, 1845–1851.

[GR73] R. Gordon and J. C. Robson, Krull dimension, Memoirs of the American Mathematical Society, No. 133, *American Mathematical Society, Providence, R.I.*, 1973, ii+78 pp.

[Her97] I. Herzog, The Ziegler spectrum of a locally coherent Grothendieck category, *Proc. London Math. Soc. (3)* **74** (1997), no. 3, 503–558.

[Kan12] R. Kanda, Classifying Serre subcategories via atom spectrum, *Adv. Math.* **231** (2012), no. 3–4, 1572–1588.

[Kan13] R. Kanda, Specialization orders on atom spectra of Grothendieck categories, arXiv:1308.3928v2, 39 pp.

[Kan14] R. Kanda, Classification of categorical subspaces of locally noetherian schemes, arXiv:1405.4473v1, 47 pp.

[Kan15] R. Kanda, Extension groups between atoms and objects in locally noetherian Grothendieck category, *J. Algebra* **422** (2015), 53–77.

[Kra97] H. Krause, The spectrum of a locally coherent category, *J. Pure Appl. Algebra* **114** (1997), no. 3, 259–271.

[Mew72] A. C. Mewborn, Quasi-simple modules and weak transitivity, *Ring theory (Proc. Conf., Park City, Utah, 1971)*, pp. 241–249, *Academic Press, New York-London*, 1972.

[Pap02] C. J. Pappacena, The injective spectrum of a noncommutative space, *J. Algebra* **250** (2002), no. 2, 559–602.

[Pop73] N. Popescu, Abelian categories with applications to rings and modules, London Mathematical Society Monographs, No. 3, *Academic Press, London-New York*, 1973, xii+467 pp.

[Ros95] A. L. Rosenberg, Noncommutative algebraic geometry and representations of quantized algebras, Mathematics and its Applications, 330, *Kluwer Academic Publishers Group, Dordrecht*, 1995, xii+315 pp.

[Sto72] H. H. Storrer, On Goldman's primary decomposition, *Lectures on rings and modules (Tulane Univ. Ring and Operator Theory Year, 1970–1971, Vol. I)*, pp. 617–661, Lecture Notes in Math., Vol. 246, *Springer-Verlag, Berlin-New York*, 1972.

[Wu88] Q. S. Wu, On an open problem of Albu and Năstăsescu, *Kexue Tongbao (English Ed.)* **33** (1988), no. 20, 1667–1668.

Graduate School of Mathematics, Nagoya University, Furo-cho, Chikusa-ku, Nagoya-shi, Aichi-ken, 464-8602, Japan
E-mail address: `kanda.ryo@a.mbox.nagoya-u.ac.jp`